

ASPIRIN WARNING

Aspirin and aspirin-containing products may cause prolonged or excessive bleeding during surgery. If you take aspirin on a regular basis to prevent stroke or heart attack, for arthritis pain, headache, fever or any other reason, discuss this with your doctor. Please be aware that many cold, cough and pain remedies contain aspirin. The following is a partial list of these products.

Alka-Seltzer*	CAMA Arthritis Pain Relief	Pain reliever tablets
Anacin*	Children's chew, Aspirin	Percodan*
Arthritis Pain Formula	Cope	Presalin
Arthritis Strength BC Powder	Damason-P	Persistin
Arthritis Strength Bufferin	Dasin	Salatim Capsules
ASA Enseals	Darvon Compound	Saleto
ASA Suppositories	Epromate	Salocol
Ascriptin	Equagesic Tablets	Sine-Off Sinus*
Ascriptin ND Caplets	Equazine M	Soma Compound*
Aspcrgum	ES Bufferin	St. Joseph's Aspirin
Aspermin	Gensan	Stanback Powder/Max
Aspertab	Goody's Headache Powder	Supac
Axotal	Isollyl Improved	Synalgos-DC Capsules
Azdonc	Lortab ASA tablets	Talwin
Bayer*	Magnaprin*	Tri-Pain Caplets
BC Powder*	Marnol	Trigesic
Buffaprin	Micrainin	Ursinus Inlay Tabs
Buaffasal	Midol	Valesin
Bufferin*	Measurin	Vanquish
Buffets	Momentum	Verin
Buffex	Norgesic	Wesprin Buffered
Buffinol	Norwich Aspirin*	Zorprin
Butalbital Compound	Pacanalgesic tablets	

*All Products

PLEASE READ ALL MEDICATION LABELS CAREFULLY. MENTION OF ASA, ASPIRIN, OR ACETYLASALICYLIC ACID MEANS THE PRODUCT CONTAINS ASPIRIN. IF YOU HAVE USED ANY ASPIRIN PRODUCTS DURING THE 7-10 DAYS PRIOR TO YOUR SURGERY, INFORM YOUR DOCTOR AND SURGEON.